

Severe New Limits on the Host Galaxies of Gamma Ray Bursts

Bradley E. Schaefer

Yale University, Physics Department, JWG 463, New Haven CT 06520-8121, USA

Received _____; accepted _____

¹schaefer@grb2.physics.yale.edu

ABSTRACT

The nature of Gamma Ray Bursts (GRBs) remains a complete mystery, despite the recent breakthrough discovery of low energy counterparts, although it is now generally believed that at least most GRBs are at cosmological distances. Virtually all proposed cosmological models require bursters to reside in ordinary galaxies. This can be tested by looking inside the smallest GRB error boxes to see if ordinary galaxies appear at the expected brightness levels. This letter reports on an analysis of the contents of 26 of the smallest regions, many from the brightest bursts. These events will have $z < 0.4$ and small uncertainties about luminosity functions, K corrections and galaxy evolutions; whereas the recent events with optical transients are much fainter and hence have high redshifts and grave difficulties in interpretation. This analysis strongly rejects the many models with peak luminosities of $10^{57} \text{photons} \cdot \text{s}^{-1}$ as deduced from the $\text{Log}N - \text{Log}P$ curve with no evolution. Indeed, the lower limit on acceptable luminosities is $6 \times 10^{58} \text{photons} \cdot \text{s}^{-1}$. The only possible solution is to either place GRBs at unexpectedly large distances (with $z > 5.9$ for the faint BATSE bursts) or to require bursters to be far outside any normal host galaxy.

Subject headings: gamma-rays: bursts

1. Introduction

The discovery of low energy counterparts for Gamma Ray Bursts (Costa et al. 1997; van Paradijs et al. 1997; Frail et al. 1997; Metzger et al. 1997) has not yet solved the problem of the location of the burster sites. Measured redshifts associated with optical transients have values $0.83 < z < 2.1$ (Metzger et al. 1997), $z = 3.42$ (Kulkarni et al. 1998), and $z = 0.0085$ (Galama et al. 1998), $z = 0.967$ (Djorgovski et al. 1998) for bursts of faint peak flux. Only in the first and last cases are the connections between the spectrum and the burst firmly resolved. The early models of cosmological bursts placed them at distances corresponding to luminosities of roughly $10^{57} \text{photons} \cdot \text{s}^{-1}$ (e.g. Fenimore et al. 1993), while it was later realized that the luminosity could be as high as $2 \times 10^{58} \text{photons} \cdot \text{s}^{-1}$ if evolutionary changes are allowed (Horváth, Mészáros, & Mészáros 1996, Totani 1997, Wijers et al. 1998). The candidate host galaxy for GRB 971214 has a very high redshift $z = 3.42$ (Kulkarni et al. 1998) which suggests that the luminosity could be even as high as $3 \times 10^{59} \text{photons} \cdot \text{s}^{-1}$. With nearby, moderate, and extreme distances all indicated, it is clear that the distance scale for cosmological bursts is not well established.

Almost all proposed burst models place GRBs inside normal galaxies (eg. Nemiroff 1994). These models can be directly tested by looking inside the smallest GRB error regions for the presence of any plausible host galaxy. The recent accurate optical transient positions are much smaller than the classical triangulation positions, but they suffer from the faintness of the burst and hence the faintness of the expected host, so that the old bright bursts are actually more restrictive for the presence of host galaxies. Indeed, the old bright bursts are at low redshifts where uncertainties in cosmology and galaxy properties are minimal, while the faint burst with optical transients are at high redshifts, where K-corrections, cosmology, luminosity functions, and evolution have large uncertainties. Nevertheless, the striking result from both old and new GRB positions is the stark absence

of galaxies at the brightness levels commonly expected.

This basic no-host-galaxy dilemma was first posed by Schaefer (1992) with improvements in analysis by Fenimore et al. (1993) and Woods & Loeb (1995). The problem is that the brightest burst regions (with the smallest areas) should typically reveal normal galaxies at around sixteenth magnitude for the usual distance scales, whereas many of these boxes are empty of galaxies to fainter than twentieth magnitude. Here, a normal galaxy is taken as one drawn randomly from the Schechter luminosity function (Binggelli, Sandage, & Tammann 1988) while the usual distance scale (Fenimore et al. 1993; Horváth, Mészáros, & Mészáros 1996) has a peak luminosity of $6 \times 10^{50} \text{ergs}^{-1}$ or $10^{57} \text{photon} \cdot \text{s}^{-1}$. Band & Hartmann (1998) introduced a Bayesian analysis procedure and concluded that an infrared data base (Larson & McLean 1997) contained no useful limits while four error boxes observed with the Hubble Space Telescope (Schaefer et al. 1997) presented a serious no-host-galaxy problem.

The previous analysis was based on samples of GRB regions either with limits only for the brightest star or galaxy in the field (Schaefer 1992; Woods & Loeb 1995) or with only four regions (Schaefer et al. 1997) or with relatively large error boxes for relatively faint bursts (Larson & McLean 1997). Schaefer et al. (1998) have accumulated a large data base of observations and has placed conservative limits on the U, B, V, R, I, J, H, and K magnitudes of the brightest possible galaxies in each of 26 of the smallest GRB error boxes. This compilation solves the limitations imposed by previous samples and allows for severe new constraints on any host galaxies. Table 1 presents these magnitudes on the brightest possible galaxy in each field along with some basic properties of the burst (Schaefer et al. 1998). The peak fluxes are for a 0.25 s from 50-300 keV in units of $\text{photons} \cdot \text{s}^{-1}$, with almost all provided by E. E. Fenimore in a recent private communication. The magnitudes should be treated as limits since some of the values represent detection thresholds and since

the host might not be the brightest galaxy in a region. The magnitudes have been corrected for extinction in our Galaxy (Zombeck 1990, Blaes et al. 1997).

Are the observed limits consistent with normal foreground galaxies with no causal connection to the GRB? The incidence of chance galaxies will depend only on the area of the error box, the galactic latitude, and the filter. Observed galaxy number densities as a function of magnitude (Jones et al. 1991; Smail et al. 1995; Gardner, Cowie, & Wainscot 1993) can be used to calculate the magnitude of the brightest expected foreground galaxy. In Table 2, the V-band limits on galaxies can be compared to the magnitudes for the brightest expected foreground galaxy (V_{beg}), with a large expected scatter due to the randomness of the brightest galaxy and the fact that some of the measures are merely limits on the brightest galaxy. The median of the differences is 0.28 ± 0.35 mag for the V-band and 0.07 ± 0.21 mag for all bands. Thus, the contents of the 26 GRB error boxes are fully consistent with chance foreground galaxies alone.

2. Analysis

A detailed analysis can place limits on the absolute magnitude of any host galaxy in each region, for some assumed peak luminosity. Specifically, the observed peak flux can be combined with the assumed peak luminosity to yield a luminosity distance to the burster and to its host galaxy. This luminosity distance can then be combined with the observed limits on the host magnitude to yield a limit on the absolute magnitude for the host. For the ensemble of boxes, the limits on the absolute magnitudes can then be compared to that expected for a normal Schechter luminosity function. The assumed peak luminosity can then be varied until agreement is reached.

Two classes of distance scales have been widely considered. The first can be called

the “no evolution” scenario, where the distances are those whose luminosity corresponds to $6 \times 10^{50} \text{ergs} \cdot \text{s}^{-1}$ (30-2000 keV) or $10^{57} \text{photons} \cdot \text{s}^{-1}$ (50-300 keV) as derived from the $\text{Log}N - \text{Log}P$ curve (Fenimore et al. 1993; Horváth, Mészáros, & Mészáros 1996), energetics limits for compact objects, and time dilation (Deng & Schaefer 1998). Almost all published papers with specific cosmological burst models require and assume this distance scale (e.g. Woosley 1993; Usov 1992; Ma & Xie 1996; Lipunov et al. 1995; Holdom & Malaney 1994; Mészáros & Rees 1993). Alternatively, an “evolutionary” distance scale might have the GRB number density following the rate of massive star formation (Totani 1997; Bagot, Zwart, & Yungelson 1998, Wijers et al. 1998), with luminosities roughly twenty times larger. This distance scale is supported by the recent possible association of a faint GRB with a $z = 3.42$ source (Kulkarni et al. 1998).

The luminosity distance is $D = (L/4\pi P)^{0.5}$, with L the peak luminosity and P the peak flux. The limit on the host’s absolute magnitude is $M = m - 5 \cdot \text{Log}(D) + 5$, where D is expressed in parsecs and m is the limit on the apparent magnitude for any host. The standard luminosity-weighted Schechter luminosity function (Binggelli, Sandage, & Tammann 1988) is adopted with $a = -1$, $M^* = -21.0$ (in the V-band for a Hubble constant of $65 \text{km} \cdot \text{s}^{-1} \cdot \text{Mpc}^{-1}$), and a low luminosity cutoff at $M = -14$.

The parameter F is the fraction of the galaxy luminosity function which is fainter than the observed limit. The most critical F measures are for bright bursts with small boxes, as these should have bright hosts and few foreground interlopers. We have data from up to seven bands for individual bursts, with some being more restrictive while others are less restrictive. Since all these restrictions simultaneously apply, we can select the minimum value, F_{min} , as providing the overall limit on the position of the galaxy within the Schechter luminosity function. This selection avoids any penalty associated with including a limit of poor sensitivity in some band.

We can quantitatively allow for the varying importance of large-versus-small boxes and faint-versus-bright bursts by forming a weighted average of the individual F_{min} values. The weight, W , will be the probability that the brightest galaxy in the field is the host and not some foreground galaxy. This probability is calculated from the magnitude of the brightest expected foreground galaxy and its position in the luminosity function for the assumed burst luminosity. The W value does not depend in any way on observations of the contents of the region. The uncertainty in $\langle F_{min} \rangle$ will be $[(\langle F_{min}^2 \rangle - \langle F_{min} \rangle^2) / \sum W]^{0.5}$.

The weighted average $\langle F_{min} \rangle$ will (for an assumed peak luminosity) be a measured statistic for comparison with models. The model $\langle F_{min} \rangle$ statistic will depend on the existence of hosts. If normal hosts are the brightest galaxy in each of the fields, then the F_{min} values will be uniformly distributed from zero to one, such that the average of all 26 values should be 0.5. With random foreground galaxies, the observed limits on the individual F_{min} values will be larger, so that $\langle F_{min} \rangle$ can only be greater than a half. Similarly, for regions where a detection threshold is reported, the individual F_{min} values can only increase. Thus, the existence of normal host galaxies requires $\langle F_{min} \rangle$ to be greater than or equal to 0.5. If hosts are not present in the error boxes, then the model $\langle F_{min} \rangle$ value can vary from near zero for low L (such that GRBs are relatively nearby and the lack of hosts is apparent) to near unity for high L (such that GRBs are very distant and the lack of hosts is not apparent against the foreground galaxies). So any acceptable model of cosmological GRBs in hosts must adopt a luminosity such that $\langle F_{min} \rangle$ is ≥ 0.5 .

The analysis must incorporate the effects of the red shift on the observed brightness of the burst and of the host galaxy. K-corrections plus E-corrections for a distribution of galaxy types have been adopted from evolutionary synthetic spectral models (Rocca-Volmerange & Guideroni 1988; Pozetti, Bruzual, & Zamorani 1996). K-corrections for the bursts have been calculated following equations 1, 2, and 4 in Fenimore et al. (1992). I have adopted

an average spectral slope index of -1.5 (see Figure 46 of Schaefer et al. 1994), although this value is varied as described below. The use of a power law spectral model is acceptable, since the $\langle F_{min} \rangle$ value is insensitive to large changes in the slope (cf. Fenimore & Bloom 1995). This procedure corrects the observed peak fluxes from 50 to 300 KeV for the effect of redshift. The Hubble Constant and the deceleration parameter enter the problem for the value of M^* as well as to calculate the E-corrections from the luminosity distance. I have adopted $H_0 = 65 km \cdot s^{-1} \cdot Mpc^{-1}$ and $q_0 = 0.5$, although these values have also been varied.

3. No Evolution Case

With these generalizations, we can first address the “no evolution” peak luminosity of $10^{57} photon \cdot s^{-1}$ ($\sim 6 \times 10^{50} ergs \cdot s^{-1}$). Table 2 presents the derived red shift (z), the most restrictive band (MRB), F_{min} , and W for all 26 bursts. The weighted average $\langle F_{min} \rangle$ is 0.141 ± 0.053 .

How robust is this result? (1) The $\langle F_{min} \rangle$ value varies from 0.153 to 0.141 as the deceleration parameter varies from 0.1 to 0.5. As the assumed Hubble Constant is varied from 50 to 80 $km \cdot s^{-1} \cdot Mpc^{-1}$, the $\langle F_{min} \rangle$ value changes from 0.125 to 0.180. (2) How large a change in the luminosity function parameters is needed to satisfy the limits? The $\langle F_{min} \rangle$ value will increase to a half by either making the low luminosity slope equal -2.1 or the M^* value over 5 magnitudes fainter. The low luminosity cutoff is unimportant for a luminosity-weighted function. (3) If only the V-band data are considered, the $\langle F_{min} \rangle$ value is 0.231 ± 0.063 . (4) If there were some [totally unsuspected] systematic error that brightened the limits in Table 1, any such errors would have to average 2.5 magnitudes to get $\langle F_{min} \rangle$ greater than 0.5. (5) If the three 1997 bursts are arbitrarily ignored, then the $\langle F_{min} \rangle$ value is 0.133 ± 0.051 . The inclusion of GRB971214 into Table 1 (with $R > 25.6$, $P_{256} \sim 2ph \cdot s^{-1}$ and $z = 3.42$) changes the $\langle F_{min} \rangle$ slightly to 0.155 ± 0.055 .

(6) If host galaxies only occupy some fraction of the error boxes, then $\langle F_{min} \rangle$ will vary linearly between the host+foreground level of 0.55 and the foreground alone level of 0.19. For an observed $\langle F_{min} \rangle$ of 0.141 ± 0.053 , the two-sigma acceptable value can be modeled by requiring that $> 84\%$ of the boxes do not have hosts. (7) If no K-corrections for the host galaxy are used, then $\langle F_{min} \rangle = 0.136 \pm 0.054$. As the average GRB spectral slope index changes from 1.0 to 2.5, the $\langle F_{min} \rangle$ value varies over a range of amplitude 0.022. (8) The analysis never uses a burst rate density so the result is independent of any assumptions on the rate density evolution.

What about the possibility of a luminosity function for the bursts? The effect on the $\langle F_{min} \rangle$ statistic will be to average it over the assumed luminosity function. To get an expected $\langle F_{min} \rangle$ greater than a half, the majority of the bursts must greatly exceed the luminosity. So a GRB luminosity function cannot solve the basic no-host-galaxy dilemma.

Let me state the no-host-galaxy dilemma in five ways with increasing generality: (1) The smallest classical GRB box is for GRB790406 with $P = 45 \text{ photons}^{-1} \cdot \text{cm}^{-2}$, so that $z = 0.09$ for the “no evolution” luminosity and an M^* galaxy should appear as $B = 17.8$ mag. Yet the region is empty of galaxies to $B = 24.29$ mag, so that any host must be > 6.5 mag fainter than M^* . (2) The existing limits on the hosts for GRB970228 and GRB970508 require the hosts to be in the bottom 0.3% and the bottom 2.2% (see F_{min} values in Table 2). For such faint galaxies, the luminosity function is not well known, yet it is well known enough to realize that both hosts are improbably faint were they to be normal galaxies. (3) For the larger sample of GRBs with $W > 0.9$, the average F_{min} values are very low, with nine of the fourteen events whose hosts must be in the bottom 4% of the luminosity function. (4) The brightest galaxy in the 26 regions has a median difference from the brightest expected foreground galaxy brightness by 0.07 ± 0.21 mag, showing that the contents of the error boxes are entirely consistent with random foreground galaxies. (5)

The $\langle F_{min} \rangle$ value is 0.141 ± 0.053 and there is no plausible means to make it ≥ 0.5 .

4. Evolutionary Case

What about the possibility that the peak luminosity is substantially brighter than the “no-evolution” value? Just such a case is expected if the GRB number density follows the rate of massive star formation (Totani 1997). The combined BATSE and PVO $LogN - LogP$ curve can be made consistent with average L values up to $2 \times 10^{58} photon \cdot s^{-1}$ ($\sim 10^{52} ergs \cdot s^{-1}$) for a careful choice of density evolution and luminosity function (Horváth, Mészáros, & Mészáros 1996). An equivalent way to quantify this limit is with the red shift of the BATSE 90% efficiency threshold ($z_{0.85}$ for $P_{256} = 0.85 photon \cdot s^{-1}$), with values ranging from two to three (Totani 1997). This luminosity from the evolutionary scenario can be tested against the limits on host galaxies.

Table 2 presents the values of z , MRB , F_{min} , and W for all 26 bursts on the assumption that $L = 2 \times 10^{58} photon \cdot s^{-1}$ (with $z_{0.85} = 3.2$). The $\langle F_{min} \rangle$ value is 0.291 ± 0.118 (versus 0.55 expected for normal host+foreground), with most of the information coming from five bursts with small boxes. All but one of the bursts have z around a half, so that K- corrections and luminosity functions are still known with some confidence. The $\langle F_{min} \rangle$ is inconsistent with the presence of host galaxies in the GRB regions at the 2.2-sigma confidence level. For $\langle F_{min} \rangle$ to be greater than a half, L must be greater than $10^{59} photons^{-1}$ (with $z_{0.85} = 7.9$), although a luminosity of $6 \times 10^{58} photon \cdot s^{-1}$ (with $z_{0.85} = 5.9$) is at the one-sigma limit. The addition of GRB971214 only slightly changes $\langle F_{min} \rangle$ to 0.323 ± 0.106 . This represents a conservative limit since (1) the brightest galaxy in the box might not be the host, (2) half the relevant magnitude limits merely represent detection thresholds, and (3) the host+foreground case predicts $\langle F_{min} \rangle$ equal to 0.55.

Uncertainties rise as the hosts are pushed to farther distances. For example, the effects of uncertainties in the cosmological parameters increase, the role of dust in obscuring young galaxies could perhaps become important, and the luminosity function might change significantly. Fortunately, the bursts used in this study are very bright (the median for bursts with $W > 0.1$ is $P_{256} = 45 \text{photons} \cdot \text{cm}^{-1} \cdot \text{s}^{-1}$) and hence close, and so cosmological uncertainties are small, there is no abnormal dust obscuration, and the luminosity function is substantially unchanged (Ellis et al. 1996). In contrast, the bursts with transients are systematically fainter (the median for the nine bursts is $P_{256} = 3.3 \text{photons} \cdot \text{cm}^{-1} \cdot \text{s}^{-1}$) and hence farther away by a factor of ~ 3 , and so have many more problems caused by cosmology, dust and evolution. This crucial difference is why host galaxy limits from the bright bursts are more constricting than limits from the faint bursts with small boxes.

5. Possible Solutions

If GRBs are cosmological, then there must be some solution to the no-host-galaxy dilemma. I can only think of two classes of solutions, first where the bursts are placed at very large cosmological distances and second where bursters do not reside in normal hosts galaxies.

If GRBs are placed at extreme distances, then the required peak luminosity is $L > 6 \times 10^{58} \text{photon} \cdot \text{s}^{-1}$ with the BATSE faint bursts at $z_{0.85} > 5.9$. Any such model would have to fine tune the cosmology and density evolution to produce the long -3/2 slope region of the PVO $\text{Log}N - \text{Log}P$ curve (Fenimore et al. 1993). Any such model would have trouble explaining the $z < 2.1$ limit for GRB970508 (Metzger et al. 1997) and the $z = 0.967$ redshift for GRB980703 (Djorgovski et al. 1998). Any such model is inconsistent (Deng & Schaefer 1998) with the observed time dilation of burst light curves (Norris et al. 1994, in't Zand & Fenimore 1996, Deng & Schaefer 1998). Finally, any such model places bursts at

distances already rejected by limits on gravitational lensing (Marani 1998).

GRBs might not reside in normal galaxies for various reasons. It might be that bursters were ejected from their galaxy of origin at high velocity so as to now appear far away. But any ejection mechanism must be $> 84\%$ efficient. Also, an analysis of six high-latitude bright bursts with small boxes shows that the area around the box is empty, forcing the average ejection-to-burst time to be $> 2 \times 10^9$ years for ejection velocities of $500 km \cdot s^{-1}$ for the canonical peak luminosity. A second alternative is that GRBs occur with equal probability per galaxy regardless of mass. Yet such a possibility is formally rejected (with $\langle F_{min} \rangle = 0.45$) while even models involving compact objects in galactic centers still have burst frequency being mass dependent. A third possibility is that the hosts are of a greatly subluminal population, with a luminosity function that has M^* fainter by 4.9 mag. Such an ad hoc assumption would require identifying an appropriate population and explaining why normal galaxies do not produce bursts. A final alternative, is that GRBs are indeed in intergalactic space, yet then there is no known source of compact objects of the required energy.

In conclusion, Gamma Ray Bursters are strongly shown to not reside in normal host galaxies at either the “no evolution” distance scale ($L = 10^{57} photon \cdot s^{-1}$ and $z_{0.85} = 0.69$) or the “evolutionary” distance scale associated with bursts as tracers of star formation ($L = 2 \times 10^{58} photon \cdot s^{-1}$ and $z_{0.85} = 3.2$). This no-host-galaxy dilemma rejects many models, and forces GRBs to either be at very large distances ($L > 6 \times 10^{58} photon \cdot s^{-1}$ with the BATSE faint bursts at $z > 5.9$) or to not be in normal host galaxies.

REFERENCES

- Bagot, P., Zwart, S. F. P. & Yungelson, L. R. 1998, A&A 332, L57
- Band, D. L. & Hartmann, D. H. 1998, ApJ 493, 555
- Binggelli, B., Sandage, A. & Tammann, G. A. 1988, ARA &A 26, 509
- Blaes, O. et al. 1997, ApJ 479, 868
- Costa, E. et al. 1997, Nature 387, 783
- Deng, M. & Schaefer, B. E. 1998, ApJ, 502, L109
- Djorgovski, S. G. et al. 1998, GCN notice 139
- Ellis, R. S. et al. 1996, MNRAS, 280, 235
- Fenimore, E. E. et al. 1992, Nature 357, 140
- Fenimore, E. E. et al. 1993, Nature 366, 40
- Fenimore, E. E. & Bloom, J. S. 1995, ApJ 453, 25
- Frail, D. et al. 1997, Nature 389, 261
- Gardner, J. P., Cowie, L. L. & Wainscoat, R. J. 1993, ApJ 415, L9
- Galama, T. J. et al. 1998, Nature, in press
- Holdom, B. & Malaney, R. A. 1994, ApJ 420, L53
- Horváth, I., Mészáros P., & Mészáros, A., 1996, ApJ 470, 56
- in’t Zand, J. J. M. & Fenimore, E. E., 1996, ApJ, 473, 622
- Jones, L. R. et al. 1991, MNRAS 249, 481

- Kulkarni, S. R. et al. 1998, *Nature* 393, 35
- Larson, S. B. & McLean, I. S. 1997, *ApJ* 491, 93
- Lipunov, V. M. et al. 1995, *ApJ* 454, 593
- Ma, F. & Xie, B. 1996, *ApJ* 462, L63
- Marani, G. F. 1998, Ph.D. Thesis, George Mason University
- Mészáros, P. & Rees, M. J. 1993, *ApJ* 405, 278
- Metzger, M. R. et al. 1997, *Nature* 387, 878
- Nemiroff, R. J. 1994, *Comments Astrophys.* 17, 189
- Norris, J. P., et al. 1994, *ApJ*, 424, 540
- Pozzetti, L., Bruzual, G. & Zamorani, G. 1996, *MNRAS* 281, 953
- Rocca-Volmerange, B. & Guideroni, B. 1988, *A&A Supp* 75, 93
- Schaefer, B. E. 1992, in *Gamma-ray Bursts: Observations, Analysis, and Theories*, eds C. Ho, R. Epstein, & E. Fenimore, (Cambridge: Cambridge Univ. Press), 107
- Schaefer, B. E. et al. 1994, *ApJSupp* 92, 285
- Schaefer, B. E. et al. 1997, *ApJ* 489, 693
- Schaefer, B. E. et al. 1998, *ApJ*, in press
- Smail, I. et al. 1995, *ApJ* 449, L105
- Totani, T. 1997, *ApJ* 486, L71
- Usov, V. V. 1992, *Nature* 357, 472

van Paradijs, J. et al. 1997, *Nature* 386, 686

Wijers, R. A. M. J., Bloom, J. S., Bagla, J. S., & Natarajan, P., 1998, *MNRAS*, 294, L13

Woods, E. & Loeb, A. 1995, *ApJ* 453, 583

Woosley, S. E. 1993, *ApJ* 405, 273

Zombeck, M. 1990, *Handbook of Space Astronomy & Astrophysics* (Cambridge: Cambridge Univ. Press)

GRB	b_{II}	Area	P_{256}	Apparent Magnitude Limits (Extinction Corrected)							
	$^{\circ}$	sq'	$ph \cdot cm^{-2} \cdot s^{-1}$	U	B	V	R	I	J	H	K
781104	-26	14	69	16.51	15.94	15.02	...	14.04	...	...	...
781119	-84	8	63	19.34	20.16	19.68	18.20	...	...	...	...
781124	80	48	99	20.90	20.62	18.69	...	17.81	...	...	...
790113	-19	78	105	...	18.44	17.18	16.03	14.53	...	...	...
790307	14	10	57	...	18.91	18.40	...	17.93	...	...	...
790313	-25	24	39	...	17.85	17.20	...	16.50	...	...	...
790325	22	2	29	22.74	22.11	20.69	19.46	...	...	...	13.56
790329	57	41	19	20.37	19.66	18.19	...	16.72	...	...	...
790331	-6	20	48	...	...	13.70	13.53	13.05	...	...	...
790406	-61	0.26	45	22.46	24.29	23.26	22.22	21.53	17.49	...	18.30
790418	-16	2.9	46	...	...	21.71	...	20.00	18.98	18.21	17.53
790613	38	0.76	15	20.92	21.06	20.17	19.45	19.33	...	...	13.58
791105	-53	35	21	20.16	20.08	19.27	...	19.31	...	...	...
791116	-75	3.7	69	22.11	22.55	21.12	19.99	19.10	...	...	...
910122	-31	19.3	42	...	20.58	19.33	...	17.95	16.86	...	15.28
910219	55	7.29	32	...	...	19.81	19.08	18.32	16.99	16.39	15.59
911118	36	12.2	41	20.06	20.17	18.89	18.31	17.60	...	...	...
920325	-44	2.1	25	...	20.39	20.42	...	...	...	...	16.99
920406	-28	0.44	73	22.27	23.42	21.99	21.84	21.39	19.35	...	18.27
920501	1	0.89	48	...	6.41	...	14.09	...	16.57	...	15.25
920711	27	1.4	23	...	20.80	20.38	20.33	...	19.05	18.16	17.47
920720	81	1.3	43	...	18.54	18.04	...	...	16.40	15.90	15.40
920723	8	4.46	95	...	...	16.88	...	16.30	17.82	...	16.04
970228	-18	0.003	10	...	...	24.63	...	24.19	...	...	...
970402	-9	2.2	0.5	18.81	18.11	17.41	17.05	16.79	...	...	...
970508	26	0.003	1.2	...	26.36	25.56	25.37	24.01	...	20.25	...

			$L = 10^{57} photon \cdot s^{-1}$				$L = 2 \times 10^{58} photon \cdot s^{-1}$			
GRB	V	V_{bfg}	z	MRB	F_{min}	W	z	MRB	F_{min}	W
781104	15.02	18.93	0.08	U	0.985	0.86	0.34	U	1.000	0.00
781119	19.68	19.47	0.08	V	0.075	0.91	0.36	V	0.888	0.07
781124	18.69	17.74	0.06	B	0.039	0.77	0.29	U	0.716	0.00
790113	17.18	17.28	0.06	B	0.249	0.68	0.28	B	1.000	0.00
790307	18.40	19.26	0.08	I	0.230	0.63	0.38	I	0.996	0.00
790313	17.20	18.41	0.10	V	0.748	0.19	0.45	I	1.000	0.00
790325	20.69	20.81	0.12	U	0.033	0.93	0.53	U	0.515	0.19
790329	18.19	17.90	0.14	U	0.366	0.25	0.65	U	1.000	0.00
790331	13.70	18.59	0.09	V	1.000	0.25	0.41	V	1.000	0.00
790406	23.26	22.78	0.09	B	0.002	1.00	0.42	B	0.107	0.82
790418	21.71	20.45	0.09	V	0.017	0.95	0.42	J	0.315	0.16
790613	20.17	21.75	0.17	B	0.212	0.94	0.73	U	0.993	0.07
791105	19.27	18.05	0.14	I	0.180	0.01	0.62	I	0.978	0.00
791116	21.12	20.22	0.08	B	0.009	0.96	0.34	B	0.292	0.28
910122	19.33	18.62	0.10	B	0.098	0.77	0.44	J	0.944	0.00
910219	19.81	19.56	0.11	V	0.139	0.83	0.50	J	0.969	0.00
911118	18.89	19.07	0.10	B	0.139	0.83	0.44	B	0.996	0.00
920325	20.42	20.76	0.13	V	0.107	0.93	0.57	K	0.864	0.00
920406	21.99	22.27	0.07	B	0.003	1.00	0.33	B	0.139	0.83
920501	...	...	0.09	J	0.057	0.97	0.41	J	0.682	0.52
920711	20.38	21.16	0.13	J	0.036	0.90	0.59	J	0.483	0.14
920720	18.04	21.23	0.10	J	0.195	0.95	0.43	J	0.985	0.35
920723	16.88	20.04	0.06	J	0.024	0.94	0.29	J	0.422	0.28
970228	24.63	27.16	0.20	I	0.003	1.00	0.90	I	0.082	0.99
970402	17.41	20.72	0.90	U	1.000	0.00	4.32	U	1.000	0.00
970508	25.56	27.16	0.58	R	0.022	1.00	2.70	B	0.117	0.98